



## King's Research Portal

DOI:

[10.1017/S1041610209008527](https://doi.org/10.1017/S1041610209008527)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Rajkumar, A. P., Thangodurai, P., Senthilkumar, P., Goyathri, K., Prince, M., & Jacob, K. S. (2009). Nature, prevalence and factors associated with depression among the elderly in a rural south Indian community. *International Psychogeriatrics*, 21(2), 372-378. <https://doi.org/10.1017/S1041610209008527>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Nature, prevalence and factors associated with depression among the elderly in a rural south Indian community

A. P. Rajkumar,<sup>1</sup> P. Thangadurai,<sup>1</sup> P. Senthilkumar,<sup>1</sup> K. Gayathri,<sup>1</sup> M. Prince<sup>2</sup> and K. S. Jacob<sup>1</sup>

<sup>1</sup>Department of Psychiatry, Christian Medical College, Vellore, India

<sup>2</sup>Institute of Psychiatry, London, U.K.

## ABSTRACT

**Background:** Depression in old age is an important public health problem causing considerable morbidity and disability worldwide. There is a dearth of community studies from India investigating geriatric depression and its associated risk factors. This study aimed to establish the nature, prevalence and factors associated with geriatric depression in a rural south Indian community.

**Methods:** We recruited 1000 participants aged over 65 years from Kaniyambadi block, Vellore, India. We assessed their socio-demographic profile, psychiatric morbidity, cognitive functioning, anthropometrics and disability status using the following structured assessment tools: Geriatric Mental State, Community Screening Instrument for Dementia, Modified CERAD 10 word list learning task, History and Aetiology Schedule Dementia Diagnosis and Subtype, WHO Disability Assessment Scale II, and Neuropsychiatric Inventory. We adopted a case control framework to study the factors associated with geriatric depression.

**Results:** Prevalence of geriatric depression (ICD-10) within the previous one month was 12.7% (95% CI 10.64–14.76%). Low income (OR 1.78; 95% CI 1.08–2.91), experiencing hunger (OR 2.58; 95% CI 1.56–4.26), history of cardiac illnesses (OR 4.75; 95% CI 1.96–11.52), transient ischemic attack (OR 2.43; 95% CI 1.17–5.05), past head injury (OR 2.70; 95% CI 1.36–5.36) and diabetes (OR 2.33; 95% CI 1.15–4.72) increased the risk for geriatric depression after adjusting for other determinants using conditional logistic regression. Having more confidants (OR 0.13; 95% CI 0.06–0.26) was the significant protective factor. Age, female gender, cognitive impairment and disability status were not significantly associated with geriatric depression. DSM-IV diagnosis of major depression was significantly correlated with experiencing hunger, diabetes, transient ischemic attack, past head injury, more disability and less nourishment; having more friends was protective.

**Conclusions:** Geriatric depression is prevalent in rural south India. Poverty and physical ill health are risk factors for depression among elderly while good social support is protective.

**Key words:** depression, dementia, risk factors, poverty, social supports

## Introduction

Depression is associated with morbidity as well as disability among the elderly (Cole and Dendukuri, 2003). They constitute a major public health problem worldwide and their prevalence rates range between 10 and 55% (Sherina *et al.*, 2004; Chi *et al.*, 2005; Tsai *et al.*, 2005; Khattri and Nepal, 2006; Kaneko *et al.*, 2007). The long-term prognosis of geriatric depression is bleak with incomplete

recovery (Jhingan *et al.*, 2001) and higher relapse rates (Mitchell and Subramaniam, 2005). Along with the physiological and psychological changes associated with aging, changes in the associated risk factors also modify the prevalence and prognosis of geriatric depression (van't Veer-Tazelaar *et al.*, 2008). Medical co-morbidity (Jongenelis *et al.*, 2004) and cognitive impairment (Steffens and Potter, 2008) have a complex bidirectional relationship with geriatric depression.

Available epidemiological studies on geriatric depression have mostly diagnosed depression by short screening instruments. The low positive predictive value (Arthur *et al.*, 1999) as well as doubtful external validity of screening instruments

Correspondence should be addressed to: K. S. Jacob, Professor of Psychiatry, Christian Medical College, Vellore 632002, India. Phone: +91 416 228 4513; Fax: +91 416 226 2268. Email: [ksjacob@cmcvellore.ac.in](mailto:ksjacob@cmcvellore.ac.in). Received 3 Nov 2008; revision requested 3 Dec 2008; revised version received 11 Dec 2008; accepted 15 Dec 2008.

(Kørner *et al.*, 2007) may yield many false positive cases and may inflate the prevalence rates of geriatric depression in community settings. Concerns regarding the different cut-off values used to diagnose depression across various settings also exist (McGivney *et al.*, 1994; Arthur *et al.*, 1999). Many studies have not employed structured assessment of depression, cognitive function and disability (Osborn *et al.*, 2003; Sherina *et al.*, 2004; Chi *et al.*, 2005; Tsai *et al.*, 2005; Khattri and Nepal, 2006; Kaneko *et al.*, 2007). Community studies from India on depression among the elderly are sparse. The only available Indian study employing the Geriatric Depression Scale (GDS), a screening tool, in a small sample of elderly has reported prevalence of up to 45.9% (Jain and Aras, 2007). Another larger Indian study evaluating the Hindi version of the GDS studied only the depressive symptoms and not the depressive disorders (Ganguli *et al.*, 1999). We aimed to study the nature, prevalence and factors associated with geriatric depression in a rural south Indian community with an adequate sample size and with detailed as well as structured methods of assessment.

## Methods

### Study design

We employed a cross-sectional study to examine the nature and prevalence of geriatric depression in a rural community. We used a case control design framework to identify factors associated with geriatric depression. This study was part of the 10/66 Dementia Research Group population based studies. The methodology employed in this study has been reported elsewhere (Jacob *et al.*, 2007a) and is only briefly mentioned here.

### Setting

Kaniyambadi block of Vellore district is in the southern Indian state of Tamil Nadu. It is spread over 127.4 square kilometers with 85 villages housing a total population of 110,000. The Department of Community Health, Christian Medical College, Vellore has been running a community health program in this block for the past five decades. It has developed a four-tier monitoring and computerized surveillance system of the residents of Kaniyambadi block, which is updated monthly. The community health workers, who live within the various village communities, provide the detailed statistics (Joseph *et al.*, 1991). They are supervised by public health nurses and physicians.

## Recruitment of participants

Participants were identified using a computerized list and a door-to-door survey. We invited all elderly people over 65 years of age to participate in this study.

## Assessment

The following instruments were employed to assess the participants: (i) Geriatric Mental State (GMS; Copeland *et al.*, 1986), (ii) Community Screening Instrument for Dementia (CSID; Hall *et al.*, 1993; 2000), (iii) World Health Organization Disability Assessment Scale II (WHODAS II; World Health Organization, 2001), (iv) Modified CERAD 10-word list learning task (Ganguli *et al.*, 1996), (v) History and Aetiology Schedule Dementia Diagnosis and subtype (HAS-DDS; Copeland *et al.*, 2002), (vi) Neuropsychiatric Inventory, and (vii) a structured proforma to assess socio-demographic characteristics, medical history and anthropometrics.

## Translation of instruments, training of personnel and quality control procedure

The Tamil versions of the GMS (Rajkumar *et al.*, 1997) and WHODAS II (World Health Organization, 2001) have been employed in earlier studies. The translation of the other instruments followed standard procedure. The training of research staff, data collection and quality control procedures were in accordance with 10/66 norms for population-based studies (Prince *et al.*, 2003). The Institutional Review Board of Christian Medical College, Vellore, approved the study.

## Data analysis

We initially analyzed the socio-demographic data, medical history, psychopathology, cognitive profile and the disability status using descriptive statistics. We derived the psychiatric diagnoses based on the computerized algorithms developed by the 10/66 Dementia Research Group (Prince *et al.*, 2003). We converted the continuous exposure variables to categorical measures by splitting them into two categories using their median values (Ragin, 1994). We calculated the odds ratios with 95% confidence intervals for various risk factors. We performed multivariate analysis using determinants, which were significant on univariate analysis, employing conditional logistic regression with the statistical software package SPSS 16.0.




Figure 1. Flowchart of recruitment of participants.

## Results

### Participant characteristics

The details of recruitment of participants are shown in Figure 1. We recruited 1000 participants and the overall response rate was 97.75%. The socio-demographic, psychiatric and disability profiles of the participants are shown in Table 1. The majority of the participants were women, lacking formal education, widowed and from the low-income category. The average age of the participants was 72.5 (SD 5.8) years.

### Nature and prevalence of geriatric depression

The prevalence of any ICD-10 (World Health Organization, 1992) depressive episode in our sample was 12.7% (95% CI 10.64–14.76 %). Among those 127 participants who had geriatric depression, 32 had a mild depressive episode (3.2%; 95% CI 2.11–4.29%), 76 had a moderate depressive episode (7.6%; 95% CI 5.96–9.24%), and 19 had a severe depressive episode (1.9%; 95% CI 1.05–2.75%). The prevalence of major depression according to DSM-IV (American Psychiatric

Association, 1994) criteria was 5.2% (95% CI 3.82–6.58). The prevalence of suicidal ideation within the previous month was 2.1% (95% CI 1.21–2.99). The prevalence of dementia defined by DSM-IV was 0.8% and by the education-adjusted 1066 criteria was 10.6%.

### Correlates of geriatric depression

The correlates of geriatric depression are shown in Table 2. Low income, experiencing hunger within the previous one month, having diabetes, cardiac illnesses, transient ischemic attacks and head injury were correlated with geriatric depression after adjustment for the effects of age, gender, education and the other determinants using conditional logistic regression analysis. Having more confidants had a significant protective effect for depressive disorders. Female gender, age, hypertension, cognitive impairment and more disability were not significantly correlated with geriatric depression.

When we ascertained major depression according to the DSM-IV criteria, the following variables were significantly associated with geriatric

**Table 1.** Socio-demographic and psychosocial profiles of the participants (n = 1000)

VARIABLE	SAMPLE N (%)	MEAN (SD)
Gender:		
Male	454 (45.4)	
Female	546 (54.6)	
Age in years		72.54 (5.87)
Education:		
No formal education	661 (66.1)	
Primary school dropouts	195 (19.5)	
Completed primary school	116 (11.6)	
Secondary education	26 (2.6)	
Graduate	2 (0.2)	
Highest Job status:		
agricultural labourer	768 (76.8)	
Others	232 (23.2)	
Marital Status:		
Never married	5 (0.5)	
Married	481 (48.1)	
Widowed/separated	514 (51.4)	
Monthly family income (in Indian Rs.)		1522 (1513)
Having private/state health insurance	0 (0)	
Lifetime smoker	530 (53.0)	
Lifetime alcohol use	194 (19.4)	
Past history of depression	22 (2.2)	
Mild cognitive impairment	37 (3.7)	
Dementia (DSM-IV)	8 (0.8)	
WHODAS total score		28.26 (18.27)

Rs. = rupees; DSM-IV = Diagnostic and Statistical Manual of Mental Disorders, 4<sup>th</sup> edition; WHODAS = World Health Organization Disability Assessment Scale.

depression after adjustment for the effects of age, gender, education and the other determinants using conditional logistic regression analysis: experiencing hunger within the previous month (OR 2.55; 95% CI 1.24–5.24); diabetes (OR 4.15; 95% CI 1.47–11.73); transient ischemic attacks (OR 3.23; 95% CI 1.31–7.96); past head injury (OR 3.69; 95% CI 1.42–9.55); mid-arm circumference less than 24 cm (OR 4.52; 95% CI 1.72–11.89); and more disability (OR 4.36; 95% CI 1.94–9.81). Having more confidants (OR 0.12; 95% CI 0.04–0.41) was protective.

The significant correlates for recent suicidal ideation, adjusting for age, education and gender using multivariate logistic regression were: being depressed (OR 4.35; 95% CI 1.75–10.78); experiencing hunger within the previous one month (OR 2.71; 95% CI 1.06–6.92); average monthly income of the family less than Rs. 1100 (OR 2.84; 95% CI 1.01–8.00). The significant correlates for more functional disability characterized by WHODAS II total score more than 25, adjusting for age, education and gender using multivariate logistic regression were: being demented (OR 2.84; 95% CI 1.78–4.53); transient ischemic attacks (OR 4.73; 95% CI 2.18–10.26); hard of hearing (OR 2.23; 95% CI 1.67–2.97); age above 71 years

(OR 1.97; 95% CI 1.52–2.56); decreased vision (OR 1.75; 95% CI 1.27–2.40); female gender (OR 1.62; 95% CI 1.19–2.20); lack of formal education (OR 1.52; 95% CI 1.09–2.12); experiencing hunger (OR 1.51; 95% CI 1.04–2.20). A severe depressive episode was not significantly associated with more functional disability (OR 1.10; 95% CI 0.43–2.81).

## Discussion

This study examined the factors associated with depression among the elderly in a rural south Indian community. Its strengths include a low refusal rate, adequate sample size and ascertaining depression with a standardized diagnostic interview. The possibility of selection bias was avoided by recruiting all consecutive elderly people and by the low refusal rate. We attempted to minimize the recall bias by collaborating with local community health workers and by using a detailed and structured interview. Its limitation was its cross-sectional design and consequent temporal ambiguity.

Physical illnesses and poverty were significantly correlated with geriatric depression, as in other studies (Cole and Dendukuri, 2003; Osborn *et al.*, 2003; Sherina *et al.*, 2004; Tsai *et al.*, 2005; Chi

**Table 2.** Factors associated with depression in the elderly

VARIABLE	DEPRESSION PRESENT (N = 127)	DEPRESSION ABSENT (N = 873)	BIVARIATE STATISTICS		MULTIVARIATE STATISTIC <sup>a</sup>	
	N (%)	N (%)	ODDS RATIO (95% CI)	P-VALUE	ODDS RATIO (95% CI)	P-VALUE
Female gender	67 (52.8)	479 (54.9)	0.92 (0.63–1.33)	0.66	0.63 (0.38–1.04)	0.07
Age above 71 years	69 (54.3)	471 (54.0)	1.02 (0.70–1.48)	0.94	0.95 (0.61–1.46)	0.80
Illiteracy	92 (72.4)	569 (65.2)	1.40 (0.93–2.12)	0.11	1.21 (0.69–2.13)	0.51
<b>Monthly family income below Rs.1100</b>	<b>88 (69.3)</b>	<b>417 (47.8)</b>	<b>2.47 (1.65–3.68)</b>	<b>0.000</b>	<b>1.78 (1.08–2.91)</b>	<b>0.02</b>
<b>Experienced hunger within past 30 days</b>	<b>45 (35.4)</b>	<b>96 (11.0)</b>	<b>4.44 (2.92–6.77)</b>	<b>0.000</b>	<b>2.58 (1.56–4.26)</b>	<b>0.000</b>
Widowed/separated status	68 (53.5)	446 (51.1)	1.10 (0.76–1.60)	0.61	0.95 (0.53–1.69)	0.85
Living alone	23 (18.1)	97 (11.1)	<b>1.77 (1.08–2.91)</b>	<b>0.03</b>	1.43 (0.78–2.61)	0.25
Living within five miles of siblings	59 (46.5)	535 (61.3)	<b>0.55 (0.38–0.80)</b>	<b>0.002</b>	0.75 (0.49–1.15)	0.19
<b>Having four or more confidants</b>	<b>8 (6.3)</b>	<b>384 (44.0)</b>	<b>0.09 (0.04–0.18)</b>	<b>0.000</b>	<b>0.13 (0.06–0.26)</b>	<b>0.000</b>
No regular physical exercise <sup>b</sup>	78 (61.4)	432 (49.5)	<b>1.63 (1.11–2.38)</b>	<b>0.01</b>	1.14 (0.73–1.79)	0.56
History of hypertension	18 (14.2)	108 (12.4)	1.17 (0.68–2.00)	0.57	1.10 (0.56–2.14)	0.79
<b>History of Diabetes mellitus</b>	<b>16 (12.6)</b>	<b>50 (5.7)</b>	<b>2.37 (1.31–4.31)</b>	<b>0.005</b>	<b>2.33 (1.15–4.72)</b>	<b>0.02</b>
History of TB within the past five years	10 (7.9)	29 (3.3)	<b>2.49 (1.18–5.24)</b>	<b>0.02</b>	1.34 (0.55–3.26)	0.52
<b>History of transient ischemic attack</b>	<b>18 (14.2)</b>	<b>26 (3.0)</b>	<b>5.37 (2.85–10.12)</b>	<b>0.000</b>	<b>2.43 (1.17–5.05)</b>	<b>0.02</b>
<b>History of cardiac illnesses</b>	<b>12 (9.4)</b>	<b>16 (1.8)</b>	<b>5.59 (2.58–12.11)</b>	<b>0.000</b>	<b>4.75 (1.96–11.52)</b>	<b>0.001</b>
<b>History of head injury with LOC</b>	<b>19 (15.0)</b>	<b>33 (3.8)</b>	<b>4.48 (2.46–8.15)</b>	<b>0.000</b>	<b>2.70 (1.36–5.36)</b>	<b>0.005</b>
Hearing impairment	50 (39.4)	264 (30.2)	<b>1.50 (1.02–2.20)</b>	<b>0.04</b>	1.46 (0.93–2.28)	0.10
Mid-arm circumference less than 24 cm	89 (70.1)	514 (58.9)	<b>1.64 (1.09–2.45)</b>	<b>0.02</b>	1.24 (0.78–1.97)	0.37
Mild cognitive impairment	2 (1.6)	35 (4.0)	0.38 (0.09–1.61)	0.19	0.49 (0.10–2.29)	0.36
Dementia <sup>c</sup>	2 (1.6)	6 (0.7)	2.31 (0.46–11.58)	0.31	0.34 (0.05–2.27)	0.27
WHODAS total score more than 25	53 (41.7)	412 (47.2)	0.80 (0.55–1.17)	0.25	0.68 (0.42–1.11)	0.13

<sup>a</sup> Adjusted odds ratios calculated using conditional logistic regression. The following variables were included in the conditional logistic regression analysis: age, gender, education, low income, experiencing hunger, living alone, living within five miles from siblings, having more confidants, no regular physical exercise, diabetes, tuberculosis, heart problems, transient Ischemic attack, head injury, hearing impairment and smaller mid-arm circumference.

<sup>b</sup> Walking for at least 500 meters less than six times per week.

<sup>c</sup> Diagnosis based on DSM-IV criteria.

LOC = loss of consciousness.

*et al.*, 2005; Khattri and Nepal, 2006; Jain and Aras, 2007; Kaneko *et al.*, 2007). However, female gender, advancing age and cognitive impairment were not associated with geriatric depression. More surprisingly, depressive disorders did not indicate more functional disability in this population. The depressed individuals in low-income communities rarely subscribe to biomedical causal models and hold more to psychosocial as well as interpersonal explanatory models for depression (Pereira *et al.*, 2007). Such explanatory models may mitigate perceived stigma (Raguram *et al.*, 1996). The meager expectations by families of their elderly relatives may also contribute towards high tolerance of depressive symptoms and functional impairment (Jacob *et al.*, 2007b).

Elderly people report depressive symptoms when they are distressed, when they are ill or are worried about the implication of their symptoms. Stressful life events and inability to cope with psychosocial problems may also lead the elderly to mention such symptoms. Consequently, the difficulty in separating distress from depression becomes a major issue (Heath, 1999). While psychiatrists suggest that brief screening instruments can easily identify people with depression (Kessler *et al.*, 1999), most general practitioners (GPs) would argue that many of those identified are distressed (Heath, 1999). The depression seen in the community and which many GPs encounter is often viewed as a result of personal and social stress, life style choices or as a product of habitual maladaptive patterns of behavior. Consequently, the general population and primary care physicians often hold psychological and social models for depression (Jacob, 2006). Psychiatrists, with their biomedical frameworks, would on the other hand argue for disease models (Kessler *et al.*, 1999). The relationship between poverty, social isolation, physical health and mental health is complex (Kuruvilla and Jacob, 2007). However, the cross-sectional and observational nature of this study mandates the need for cohort designs to confirm associations and randomized trials to assess intervention.

### Conflict of interest

None.

### Description of authors' roles

K. S. Jacob and M. Prince designed the study and supervised data collection and analysis. P. Senthilkumar and K. Gayathri collected the data and assisted with writing the paper. A.P. Rajkumar,

P. Thangadurai and K. S. Jacob analyzed the data and wrote the paper.

### Acknowledgments

This study is part of the population-based investigations of the 10/66 Dementia Research Group of Alzheimer's Disease International. It was supported by grants from the World Health Organization and the Institute of Psychiatry, London. We thank all the study participants as well as their families and the community health workers for their cooperation, and the research fellows Ms. A. Kanagathara and Ms. C. Symon for data collection.

### References

- American Psychiatric Association** (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn (DSM- IV). Washington, DC: American Psychiatric Association.
- Arthur, A., Jagger, C., Lindesay, J., Graham, C. and Clarke, M.** (1999). Using an annual over-75 health check to screen for depression: validation of the short Geriatric Depression Scale (GDS15) within general practice. *International Journal of Geriatric Psychiatry*, 14, 431–439.
- Chi, I. et al.** (2005). Prevalence of depression and its correlates in Hong Kong's Chinese older adults. *American Journal of Geriatric Psychiatry*, 13, 409–416.
- Cole, M. G. and Dendukuri, N.** (2003) Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *American Journal of Psychiatry*, 160, 1147–1156.
- Copeland, J. R., Dewey, M. E. and Griffith-Jones, H. M.** (1986). A computerized psychiatric diagnostic system and case nomenclature for elderly subjects: GMS and AGE-CAT. *Psychological Medicine*, 16, 89–99.
- Copeland, J. R., Prince, M., Wilson, K. C., Dewey, M. E., Payne, J. and Gurland, B.** (2002). The Geriatric Mental State examination in the 21st century. *International Journal of Geriatric Psychiatry*, 17, 727–732.
- Ganguli, M. et al.** (1996). Cognitive test performance in a community-based nondemented elderly sample in rural India: the Indo-U.S. Cross-National Dementia Epidemiology Study. *International Psychogeriatrics*, 8, 507–524.
- Ganguli, M., Dube, S., Johnston, J. M., Pandav, R., Chandra, V. and Dodge, H. H.** (1999). Depressive symptoms, cognitive impairment and functional impairment in a rural elderly population in India: a Hindi version of the geriatric depression scale (GDS-H). *International Journal of Geriatric Psychiatry*, 14, 807–820.
- Hall, K. S., Hendrie, H. C. and Brittain, H. M.** (1993). The development of a dementia-screening interview in two distinct languages. *International Journal of Methods in Psychiatric Research*, 3, 1–28.
- Hall, K. S., Gao, S., Emsley, C. L., Ogunniyi, A. O., Morgan, O. and Hendrie, H. C.** (2000). Community

- screening interview for dementia (CSI 'D'): performance in five disparate study sites. *International Journal of Geriatric Psychiatry*, 15, 521–531.
- Heath, I.** (1999). Commentary: there must be limits to the medicalisation of human distress. *BMJ*, 318, 439–440.
- Jacob, K. S.** (2006). The cultures of depression. *National Medical Journal of India*, 19, 218–220.
- Jacob, K. S., Senthil Kumar, P., Gayathri, K., Abraham, S. and Prince, M. J.** (2007a). The diagnosis of dementia in the community. *International Psychogeriatrics*, 19, 669–678.
- Jacob, K. S., Senthil Kumar, P., Gayathri, K., Abraham, S. and Prince, M. J.** (2007b). Can health workers diagnose dementia in the community? *Acta Psychiatrica Scandinavica*, 116, 125–128.
- Jain, R. K. and Aras, R. Y.** (2007). Depression in geriatric population in urban slums of Mumbai. *Indian Journal of Public Health*, 51, 112–113.
- Jhingan, H. P., Sagar, R. and Pandey, R. M.** (2001). Prognosis of late-onset depression in the elderly: a study from India. *International Psychogeriatrics*, 13, 51–61.
- Jongenelis, K., Pot, A. M., Eisses, A. M., Beekman, A. T., Kluiters, H. and Ribbe, M. W.** (2004). Prevalence and risk indicators of depression in elderly nursing home patients: the AGED study. *Journal of Affective Disorders*, 83, 135–142.
- Joseph, A., Joseph, K. S. and Kamaraj, K.** (1991). Use of computers in primary health care. *International Journal of Health Sciences*, 2, 93–101.
- Kaneko, Y., Motohashi, Y., Sasaki, H. and Yamaji, M.** (2007). Prevalence of depressive symptoms and related risk factors for depressive symptoms among elderly persons living in a rural Japanese community: a cross-sectional study. *Community Mental Health Journal*, 43, 583–590.
- Kessler, D., Lloyd, K., Lewis, G. and Gray, D. P.** (1999). Cross sectional study of symptom attribution and recognition of depression and anxiety in primary care. *BMJ*, 318, 436–439.
- Khatti, J. B. and Nepal, M. K.** (2006). Study of depression among geriatric population in Nepal. *Nepal Medical College Journal*, 8, 220–223.
- Korner, A. et al.** (2007). Rating scales for depression in the elderly: external and internal validity. *Journal of Clinical Psychiatry*, 68, 384–389.
- Kuruvilla, A. and Jacob, K. S.** (2007). Poverty, social stress and mental health. *Indian Journal of Medical Research*, 126, 273–278.
- McGivney, S. A., Mulvihill, M. and Taylor, B.** (1994). Validating the GDS depression screen in the nursing home. *Journal of the American Geriatrics Society*, 42, 490–492.
- Mitchell, A. J. and Subramaniam, H.** (2005). Prognosis of depression in old age compared to middle age: a systematic review of comparative studies. *American Journal of Psychiatry*, 162, 1588–1601.
- Osborn, D. P. et al.** (2003). Factors associated with depression in a representative sample of 14,217 people aged 75 and over in the United Kingdom: results from the MRC trial of assessment and management of older people in the community. *International Journal of Geriatric Psychiatry*, 18, 623–630.
- Pereira, B., Andrew, G., Pednekar, S., Pai, R., Pelto, P. and Patel, V.** (2007). The explanatory models of depression in low-income countries: listening to women in India. *Journal of Affective Disorders*, 102, 209–218.
- Prince, M. J., Acosta, D., Chiu, H., Scazufca, M. and Varghese, M.** (2003). Dementia diagnosis in developing countries: a cross-cultural validation study. *Lancet*, 361, 909–917.
- Ragin, C.** (1994). *Constructing Social Research*. Thousand Oaks, CA: Pine Forge Press, Sage.
- Raguram, R., Weiss, M. G., Channabasavanna, S. M. and Devins, G. M.** (1996). Stigma, depression, and somatization in South India. *American Journal of Psychiatry*, 153, 1043–1049.
- Rajkumar, S., Kumar, S. and Thara, R.** (1997). Prevalence of dementia in a rural setting: a report from India. *International Journal of Geriatric Psychiatry*, 12, 702–707.
- Sherina, M. S., Rampal, L. and Mustaqim, A.** (2004). The prevalence of depression among the elderly in Sepang, Selangor. *Medical Journal of Malaysia*, 59, 45–49.
- Steffens, D. C. and Potter, G. G.** (2008). Geriatric depression and cognitive impairment. *Psychological Medicine*, 38, 163–175.
- Tsai, Y. F., Yeh, S. H. and Tsai, H. H.** (2005). Prevalence and risk factors for depressive symptoms among community-dwelling elders in Taiwan. *International Journal of Geriatric Psychiatry*, 20, 1097–1102.
- van't Veer-Tazelaar, P. J. et al.** (2008). Depression in old age (75+): the PIKO study. *Journal of Affective Disorders*, 106, 295–299.
- World Health Organization** (1992) *The ICD-10 classification of mental and behavioural disorders: Diagnostic Criteria for Research*. Geneva: WHO.
- World Health Organization** (2001). *WHODAS II Disability Assessment Schedule*. Geneva: WHO.